arXiv:1611.06568v1 [hep-ph] 20 Nov 2016

Left-right symmetric extensions of 3-3-1 models

Edison T. Franco*

Universidade Federal do Tocantins - Campus Universitário de Araguaína

Av. Paraguai (esquina com Urixamas), Cimba

Araguaína - TO, 77814-970, Brazil

V. Pleitez†

Instituto de Física Teórica–Universidade Estadual Paulista

R. Dr. Bento Teobaldo Ferraz 271, Barra Funda

São Paulo - SP, 01140-070, Brazil

(Dated: 11/20/2016)

Abstract

We propose four left-right symmetric extensions of several chiral 3-3-1 models. Although they have some common features they also have important difference due to different representation content.

PACS numbers: 12.60.Fr 12.15.-y 14.60.Pq

*Electronic address: edisonfranco@uft.edu.br

†Electronic address: vicente@ift.unesp.br

In the electroweak standard model (ESM), left-handed fermions transform in a different way than the right-handed components. Hence, it is a chiral model. In this way parity violation is accommodated but not explained. The parity issue was the motivation of the early left-right symmetric extensions of the ESM [1–3]. In this sort of models the electroweak gauge symmetry is extended to $SU(2)_L \otimes SU(2)_R \otimes U(1)_{B-L}$ [4] and, besides the explanation of parity violation as a spontaneously broken symmetry, the model are able to generate Dirac [3] or Majorana [5, 6] masses for neutrinos. A left-right symmetric extension can be implemented for all chiral models.

Here, we will consider that sort of extensions for models with 3-3-1 chiral symmetry in such a way that the gauge symmetries are $G_{3331} \equiv SU(3)_C \otimes SU(3)_L \otimes SU(3)_R \otimes U(1)_X$ with the charge operator defined as $Q = T_3^L + T_3^R - \sqrt{3}(T_8^L + T_8^R) + X$ (models Ia, Ib) and $Q = T_3^L + T_3^R - 1/\sqrt{3}(T_8^L + T_8^R) + X$ (model IIa,IIb) [7]. In all these extensions the anomalies are canceled considering three generations only as in the chiral 3-3-1 version since left- and right- fermion triplets have the same $U(1)_X$-charge. A model similar with our model IIa was proposed in Ref. [8] but those authors used only scalar triplets and for this reason fermion masses are generated only by five-dimensional operators. Here we will use in both sort of models only renormalizable interactions.

In the model Ia, the particle content is [9–11]: left-handed quarks $Q_{aL} = (d'_a, -u'_a, j'_a)^T_L \sim (\mathbf{3}, \mathbf{3}^*_L, \mathbf{1}_R, -1/3)$, $Q_{3L} = (u'_3, d'_3, J')^T_L \sim (\mathbf{3}, \mathbf{3}_L, \mathbf{1}_R, 2/3)$; and the right-handed quarks: $Q_{aR} = (d'_a, -u'_a, j'_a,)^T_R \sim (\mathbf{3}, \mathbf{1}_L, \mathbf{3}^*_R, -1/3)$, $Q_{3R} = (u'_3, J', d'_3)^T_R \sim (\mathbf{3}, \mathbf{1}_L, \mathbf{3}_R, 2/3)$. In the lepton sector we have left-handed leptons: $\psi_{lL} = (\nu'_l, l', l'^c)^T_L \sim (\mathbf{1}, \mathbf{3}_L, \mathbf{1}_R, 0)$; and right-handed leptons: $\psi_{lR} = (\nu'_l, E'^c_l, E'_l)^T_R \sim (\mathbf{1}, \mathbf{1}_L, \mathbf{3}_R, 0)$. Primed fields denote symmetry eigenstates.

The left-handed and right-handed fields are not equivalent since in each triplet the charged degrees of freedom are complete. Only neutrino has one component in each sector. At the energy scale at which the symmetry is valid all fields are uncharged and massless and this does not matter.

In the scalar sector we introduce the following multiplets: $T \sim (\mathbf{1}, \mathbf{3}_L, \mathbf{3}^*_R, 0)$, $P \sim (\mathbf{1}, \mathbf{3}_L, \mathbf{3}_R, 1)$, which couple to quarks, T also couples with leptons but the sextets, $S_L \sim (\mathbf{1}, \mathbf{6}^*_L, \mathbf{1}_R, 0)$ and $S_R \sim (\mathbf{1}, \mathbf{1}_L, \mathbf{6}^*_R, 0)$, couple only with leptons. The electric charge assign-

ment of the bi-triplet T, P and the sextets $S_{L,R}$ are as follows:

$$T = \begin{pmatrix} t_1^0 & t_1^- & t_2^+ \\ t_3^+ & t_2^0 & t_1^{++} \\ t_4^- & t_2^{--} & t_3^0 \end{pmatrix}, \quad P = \begin{pmatrix} \rho_1^+ & \rho_1^0 & \rho_1^{++} \\ \rho_2^0 & \rho_2^- & \rho_3^+ \\ \rho_2^{++} & \rho_4^+ & \rho^{+++} \end{pmatrix}, \quad S_{L,R} = \begin{pmatrix} s_{1L,R}^0 & \frac{s_{1L,R}^+}{\sqrt{2}} & \frac{s_{2L,R}^-}{\sqrt{2}} \\ \frac{s_{1L,R}^+}{\sqrt{2}} & s_{1L,R}^{++} & \frac{s_{2L,R}^0}{\sqrt{2}} \\ \frac{s_{2L,R}^-}{\sqrt{2}} & \frac{s_{2L,R}^0}{\sqrt{2}} & s_{2L,R}^{--} \end{pmatrix}, \quad (1)$$

Notice that P involves a scalar Higgs with triple charge. We define

$$\langle T \rangle = \begin{pmatrix} v_d & 0 & 0 \\ 0 & v_u & 0 \\ 0 & 0 & v_j \end{pmatrix}, \quad \langle P \rangle = \begin{pmatrix} 0 & V_d & 0 \\ V_u & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad \langle S_{L,R} \rangle = \begin{pmatrix} v_{L,R1} & 0 & 0 \\ 0 & 0 & \frac{v_{L,R2}}{\sqrt{2}} \\ 0 & \frac{v_{L,R2}}{\sqrt{2}} & 0 \end{pmatrix} \quad (2)$$

Notice that because of the bi-triplets $P, \bar{P}$ the symmetry is breakdown to $U(1)_Q$. Here we will assume, for the sake of simplicity, that all VEVs are real.

The Yukawa interactions in the quark sector are:

$$-\mathcal{L}_q^Y = \bar{Q}_{aL} G_{ab} Q_{bR} T^\dagger + \bar{Q}_{3L} G_{3a} Q_{aR} P + \bar{Q}_{aL} G_{a3} Q_{3R} P^\dagger + \bar{Q}_{3L} G_3 Q_{3R} T + H.c., \quad (3)$$

$a, b, 3$ are generation indices, and we have omitted $SU(3)$ indices and summation over flavors. Notice that with only the bi-triplet T only the first two families mix, G_{ab} is a 2×2 matrix, and G_3 is a constant. Besides, $\langle T \rangle \neq 0$ leaves an extra $U(1)$ unbroken symmetry. All of these is corrected by adding a second bi-triplet P carrying $U(1)_X$-charge.

The Yukawa interactions in the lepton sector are:

$$-\mathcal{L}_l^Y = \bar{\psi}_{iL} G_{ij}^l \psi_{jR} T + \overline{(\psi_{iL})^c} G_{ij}^S \psi_{jL} S_L + \overline{(\psi_{iR})^c} G_{ij}^S \psi_{jR} S_R] + H.c. \quad (4)$$

where summation over $i, j = e, \mu, \tau$ has been omitted.

We add a parity $\mathcal{P}$ as follows:

$$\begin{aligned} & g_L \leftrightarrow g_R,\ Q_{aL} \leftrightarrow Q_{aR},\ Q_{3L} \leftrightarrow Q_{3R},\ \psi_L \leftrightarrow \psi_R, \\ & T \leftrightarrow T^\dagger,\ P \leftrightarrow P,\ S_L \leftrightarrow S_R,\ W_L \leftrightarrow W_R\,, \end{aligned} \quad (5)$$

which implies that $G = G^\dagger$ in Eq. (3), $G^l = G^{l\dagger}$ in Eq. (4) and thus all mass matrices are Hermitian.

In the quark sector we have, for type-u quarks in the basis $(-u_1', -u_2', u_3')$ and in the type-d sector in the basis (d_1', d_2', d_3') the respective mass matrices are given by

$$M^u = \begin{pmatrix} G_{11} v_u & G_{12} v_u & G_{13} V_u \\ G_{12}^* v_u & G_{22} v_u & G_{23} V_u \\ G_{13}^* V_u & G_{23}^* V_u & G_3 v_d \end{pmatrix}, \quad M^d = \begin{pmatrix} G_{11} v_d & G_{12} v_d & G_{13} V_d \\ G_{12}^* v_d & G_{22} v_d & G_{23} V_d \\ G_{13}^* V_d & G_{23}^* V_d & G_3 v_u \end{pmatrix}. \quad (6)$$

Since the mass matrices are symmetric ($M^{u,d} = (M^{u,d})^\dagger$) they are diagonalized by one unitary matrix and not by two unitary matrices. The exotic quarks have their masses given by $M^j = Gv_j$ and $M^J = G_3 v_j$.

In the lepton sector we have that leptons with the same electric are separated in two 3×3 matrices. In the lepton sector we have

$$M^l = \begin{pmatrix} [G^S + \left(G^S\right)^T]\frac{v_{2L}}{\sqrt{2}} & G^l v_u \\ \left(G^l\right)^T v_j & [G^S + \left(G^S\right)^T]\frac{v_{2R}}{\sqrt{2}} \end{pmatrix}, \quad M^\nu = \begin{pmatrix} 2G^S v_{1L} & G^l v_d \\ \left(G^l\right)^T v_d & 2G^S v_{1R} \end{pmatrix}, \quad (7)$$

where the mass term for the neutrinos is written in the form $-\mathcal{L}_\nu^{mass} = \frac{1}{2}\overline{n_L} M^\nu \left(n_L\right)^c + H.c.$ in the basis $n_L = (\nu'_L, \nu'^c_R)^T$, where $G^{S,l}$ are arbitrary 3×3 matrices. If $v_{R2} \gg v_{L2}, v_u, v_j$ and $v_{R1} \gg v_{L1}, v_d$, the exotic leptons are heavier than the known leptons, and the right-handed neutrino are heavier than the left-handed (active) ones. In fact, in this model there is a type-I and type-II seesaw mechanisms.

In the present case both left- and right- handed neutrinos are Majorana particles but if, instead of the sextets S_L and S_R we introduces triplets, we would have Dirac neutrinos. For instance $\eta_L \sim (\mathbf{1},\mathbf{3},\mathbf{1},0), \eta_R \sim (\mathbf{1},\mathbf{1},\mathbf{3},0)$, the Yukawa interactions $G_L \psi_L^T \epsilon C \psi_L \eta_L$ and $G_R \psi_R^T \epsilon C \psi_R \eta_R$ are allowed but the masses obtained are not still realistic ones because G_L and G_R are 3×3 antisymmetric matrices.

A different model (model Ib) can be obtained with the same $\beta = -\sqrt{3}$, if l^c is substituted by a new charged lepton E^+ that may be considered as a particle instead of an antiparticle. See Ref. [12]. In this case $\psi_{lL} = (\nu'\, l', E_l^{+\prime})_L^T \sim (\mathbf{3},\mathbf{1},0)$ and $\psi_{lR} = (\nu'\, l', E_l^{+\prime})_R^T \sim (\mathbf{1},\mathbf{3},0)$, and all the lepton masses are generated by the Yukawa interactions like Eq. (4). The charged lepton mass matrix is now given by

$$M^l = \begin{pmatrix} G^l v_u & \left[G^S + \left(G^S\right)^T\right]\frac{v_{2L}}{\sqrt{2}} \\ \left[G^S + \left(G^S\right)^T\right]\frac{v_{2R}}{\sqrt{2}} & \left(G^l\right)^T v_j \end{pmatrix}, \quad (8)$$

where the basis of charged leptons is $\ell'_L = (l'_L, E'_L)^T$. The neutrino and quark sectors are the same as in the previous model. Yet, if the sextets are not introduced the neutrinos are Dirac particles and the exotic charged leptons decouple from the light ones. This case is not realistic since the mass matrices of charged leptons and neutrinos are proportional, thus inducing a trivial PMNS.

The details of the scalar potential in this model Ia is given by

$$
\begin{aligned}
V &= \mu_T^2 \mathrm{Tr} T^\dagger T + \mu_P^2 \mathrm{Tr} P^\dagger P + \mu_S^2 \mathrm{Tr}(S_L^* S_L + S_R^* S_R) + \lambda_1 \mathrm{Tr}(T^\dagger T)^2 + \lambda_2 (\mathrm{Tr}(T^\dagger T))^2 \\
&+ \lambda_3 \mathrm{Tr}(P^\dagger P)^2 + \lambda_4 (\mathrm{Tr}(P^\dagger P))^2 + \lambda_5 \mathrm{Tr}(S_L^* S_L + S_R^* S_R)^2 + \lambda_6 (\mathrm{Tr}(S_L^* S_L + S_R^* S_R))^2 \\
&+ \mathrm{Tr}[(S_L S_L^* + S_R S_R^*)(\lambda_7 T^\dagger T + \lambda_8 P^\dagger P)] + \mathrm{Tr}[(S_L^* S_L + S_R^* S_R)] \mathrm{Tr}[\lambda_9 (T^\dagger T) + \lambda_{10} (P^\dagger P)] \\
&+ \lambda_{11} [\mathrm{Tr}(T^\dagger T P^\dagger P) + \mathrm{Tr}(T T^\dagger P^\dagger P)] + \lambda_{12} \mathrm{Tr}(T^\dagger T) \mathrm{Tr}(P^\dagger P) \\
&+ \lambda_{13} [\mathrm{Tr}(P^\dagger S_L^*) \mathrm{Tr}(P S_L) + \mathrm{Tr}(P^\dagger S_R^*) \mathrm{Tr}(P S_R)] + \lambda_{14} [\mathrm{Tr}\left(T^\dagger T P^T P^*\right) + \mathrm{Tr}\left(T T^\dagger P^T P^*\right)] \\
&+ \lambda_{15} [(S_R^* T)(S_L T) + H.c.] + f_1 [(TTT) + H.c.] + f_2 [(S_L S_L S_L) + H.c.] + f_3 [(S_R S_R S_R) \\
&+ H.c.], \qquad (9)
\end{aligned}
$$

and in model Ib we omit in Eq. (9) all terms involving the sextets.

In model IIa we have the following representation content: left-handed quarks $Q_{aL} = (d'_a, -u'_a, D'_a)^T_L \sim (\mathbf{3}, \mathbf{3}^*_L, \mathbf{1}_R, 0)$, $Q_{3L} = (u'_3, d'_3, U')^T_L \sim (\mathbf{3}, \mathbf{3}_L, \mathbf{1}_R, 1/3)$; and the right-handed quarks: $Q_{aR} = (d'_a, -u', D')^T_R \sim (\mathbf{3}, \mathbf{1}_L, \mathbf{3}^*_R, 0)$, $Q_{3R} = (u'_3, d'_3, U')^T_R \sim (\mathbf{3}, \mathbf{1}_L, \mathbf{3}_R, 1/3)$. In the lepton sector we have left-handed leptons: $\psi_{lL} = (\nu'_l, l', N'_l)^T_L \sim (\mathbf{1}, \mathbf{3}_L, \mathbf{1}_R, -1/3)$; and right-handed leptons: $\psi_{lR} = (\nu'_l, l', N'_l)^T_R \sim (\mathbf{1}, \mathbf{1}_L, \mathbf{3}_R, -1/3)$. Primed fields denote symmetry eigenstates. Again, here N'_{lL} may be identified with $(\nu_{lR})^c$ as in the models of Refs. [13, 14], then N'_{lR} is a new neutral lepton and we call this model IIb; or both N'_{lL} and N'_{lR} are new neutral leptons as in the model of Refs. [15, 16]. In model II, the scalar multiplets are: $T' \sim (\mathbf{1}, \mathbf{3}_L, \mathbf{3}^*_R, 0)$, and $H \sim (\mathbf{1}, \mathbf{3}_L, \mathbf{3}_R, 1/3)$, defined as follows

$$
T' = \begin{pmatrix} \tilde{t}_1^0 & \tilde{t}_1^- & \tilde{t}_2^0 \\ \tilde{t}_2^+ & \tilde{t}_3^0 & \tilde{t}_3^+ \\ \tilde{t}_4^0 & \tilde{t}_4^- & \tilde{t}_5^0 \end{pmatrix}, \; H = \begin{pmatrix} h_1^+ & h_1^0 & h_2^+ \\ h_2^0 & h_3^- & h_3^0 \\ h_4^+ & h_4^0 & h_5^+ \end{pmatrix}. \qquad (10)
$$

We define the VEVs in the following form:

$$
\langle T' \rangle = \begin{pmatrix} v_d & 0 & v_R \\ 0 & v_u & 0 \\ v_L & 0 & v_D \end{pmatrix}, \; \langle H \rangle = \begin{pmatrix} 0 & V_u & 0 \\ V_d & 0 & V_D \\ 0 & V_U & 0 \end{pmatrix}. \qquad (11)
$$

The Yukawa interactions are given by

$$
-\mathcal{L}_q^Y = \sum_{a=1,2} [\bar{Q}_{aL} G_{ab} Q_{bR} T'^\dagger + \bar{Q}_{3L} G_{3a} Q_{aR} H + \bar{Q}_{aL} G_{a3} Q_{3R} H^\dagger] + \bar{Q}_{3L} G_3 Q_{3R} T' + H.c. \qquad (12)
$$

in the quark sector and

$$-\mathcal{L}_l^Y \;=\; \sum_{i,j=1}^{3} \overline{(\psi_{iL})} G_{ij}^l \psi_{jR} T' + H.c., \tag{13}$$

in the lepton sector.

The parity $\mathcal{P}$ in this case is

$$g_L \leftrightarrow g_R,\; Q_{aL} \leftrightarrow Q_{aR},\; Q_{3L} \leftrightarrow Q_{3R},\; \psi_L \leftrightarrow \psi_R,\; T' \leftrightarrow T'^{\dagger},\; H \leftrightarrow H,\; W_L \leftrightarrow W_R\;. \tag{14}$$

Again, all the G matrices entries obey $G_{ij} = G_{ji}^*$.

The mass matrix in the quark sector are:

$$M^u = \begin{pmatrix} v_u G_{11} & v_u G_{12} & V_u G_{13} & V_U G_{13} \\ v_u G_{12}^* & v_u G_{22} & V_u G_{23} & V_U G_{23} \\ V_u G_{13}^* & V_u G_{23}^* & v_d G_3 & v_R G_3 \\ V_U G_{13}^* & V_U G_{23}^* & v_L G_3 & v_D G_3 \end{pmatrix} \tag{15}$$

for the u-type in the basis $(-u_1', -u_2', u_3', U')$, and

$$M^d = \begin{pmatrix} v_d G_{11} & v_d G_{12} & V_d G_{13} & v_L G_{11} & v_L G_{12} \\ v_d G_{12}^* & v_d G_{22} & V_d G_{23} & v_L G_{12}^* & v_L G_{22} \\ V_d G_{13}^* & V_d G_{23}^* & v_u G_3 & V_D G_{13}^* & V_D G_{23}^* \\ v_R G_{11} & v_R G_{12} & V_D G_{13} & v_D G_{11} & v_D G_{12} \\ v_R G_{12}^* & v_R G_{22} & V_D G_{23} & v_D G_{12}^* & v_D G_{22} \end{pmatrix} \tag{16}$$

for the d-type in the basis $(d_1', d_2', d_3', D_1', D_2')$. Notice that if we set $v_{L,R} = V_{U,D} = 0$ the known quarks do not couple with the new ones U' and $D_{1,2}'$, however it is better assume only that $v_x/v \ll 1$, where $v_x = v_{L,R}, V_{U,D}$ and v any other VEV. This is because if there is no mixing among all quarks of the same charge, an automatic $\mathbb{Z}_2$ survive and the extra quarks are stable, or at least long lived, unless the interactions with the scalars mix all them other. The masses in the lepton sector are given by

$$M^l = G^l v_u, \quad M^\nu = \begin{pmatrix} 0 & M_D \\ M_D^T & 0 \end{pmatrix}, \quad M_D = \begin{pmatrix} G^l v_d & G^l v_R \\ G^l v_L & G^l v_D \end{pmatrix}, \tag{17}$$

in the basis $n_L^T = (\nu_L', N_L', \nu_R'^c, N_R'^c)^T$. In this case the light neutrinos become Dirac particles. It is clear to see from Eqs. (17) that the light neutrinos decouple the heavy ones only if

$v_D = 0$, but this case is not realistic since the PMNS matrix is also the identity matrix. Notice that we obtain the most general mass matrix for neutrino if we assume a model as those in Ref. [15, 16].

The scalar potential in this case is given by

$$\begin{aligned} V &= \mu_{T'}^2 \mathrm{Tr}\left(T'^\dagger T'\right) + \mu_H^2 \mathrm{Tr}\left(H^\dagger H\right) + \lambda_1 \mathrm{Tr}\left(T'^\dagger T'\right)^2 + \lambda_2 \left(\mathrm{Tr}\left(T'^\dagger T'\right)\right)^2 + \lambda_3 \mathrm{Tr}\left(H^\dagger H\right)^2 \\ &+ \lambda_4 \left(\mathrm{Tr}\left(H^\dagger H\right)\right)^2 + \lambda_5 [\mathrm{Tr}\left(T'^\dagger T' H^\dagger H\right) + \mathrm{Tr}\left(T' T'^\dagger H^\dagger H\right)] + \lambda_6 \mathrm{Tr}\left(T'^\dagger T'\right) \mathrm{Tr}\left(H^\dagger H\right) \\ &+ \lambda_7 [\mathrm{Tr}\left(T'^\dagger T' H^T H^*\right) + \mathrm{Tr}\left(T' T'^\dagger H^T H^*\right)] + f_1 \left(\mathrm{Tr}\left(T' T' T'\right) + H.c.\right). \end{aligned} \tag{18}$$

Details of the model IIb will be given elsewhere but advance that there are important differences with respect the model IIa, as those between model Ia and model Ib.

All 3-3-1 models have interesting new processes that deserve to be searched in accelerators. In particular the minimal model ($\beta = -\sqrt{3}$), predicts new resonances formed by exotic quarks in processes like

$$\begin{aligned} &pp(J\bar{J}u\bar{u}V^-V^+) \to e_R^- e_L^+ + \nu_L^c \nu_R (\text{missing energy}) + jets, \\ &pp(J\bar{J}d\bar{u}V^-U^{++}) \to e_L^+ e_R^- e_L^- e_R^+ + \nu_L^c (\text{missing energy}) + jets, \\ &pp(j\bar{j}d\bar{d}V^-V^+) \to e_R^+ e_L^- + \nu_L \nu_L (\text{missing energy}) + jets, \\ &pp(j\bar{j}u\bar{u}U^{--}U^{++}) \to l^- l^+ l^- l^+ + jets, \end{aligned} \tag{19}$$

which are interesting because the resonances involving the exotic quarks are long-lived, since they are protected by a $\mathbb{Z}_2$ symmetry which is broken by scalar interactions. In the left-right symmetric extensions this model has interactions with like those in Eq. (3) with the bi-triplet T and quarks: $G_{a3}\overline{J}[d'_a\rho^{++} + u'_3\rho^+ + j'_a\rho^{+++}]$, this is an example of interactions in which we can recognize that the exotic particles $J, j, \rho^{+++,++,+}$ are produced by pairs since they are odd under $\mathbb{Z}_2$ but u, d-type quarks are even. In this case the exotic particles will be stable. Fortunately, we have verified that all singly charged scalars mix in the mass matrices and that in the scalar potential there are interactions of exotic scalars and those coupled with known quarks and leptons as for instance that in the λ_{13} in Eq. (9), $\rho^{+++} s_{2L}^{--} s_{1L}^- \rho_1^{0*}(\rho_2^{0*})$, and from the Yukawa interaction in Eq. (4), we see that there are interactions like $G_{ij}^S[\overline{(l'_{iL})^c} s_{1L}^+ \nu'_{jL} + \overline{l'_{iR}} s_{2L}^{--} (l'_{jR})^c]$ allowing the decay of ρ^{+++} to leptons through s_{1L}^+ and s_{2L}^{++}. Hence, in particular in model I, there is a resonance with triple electric charge $\Delta^{+++}(\bar{j}J)$ which can decay into the scalar ρ^{+++} and, since there are interactions like those

in the λ_{13} term discussed above, the decay $\rho^{+++} \to s_1^+ s_2^{++}$ does occurs, after ρ_1^0 and/or ρ_2^0 get VEV, and $\rho^{+++} \to s_{1L}^+ s_{2L}^{++} \to l_R^+ l_R^+ l_L^+ \nu_L^c$ is allowed. Hence, the decay $\Delta^{+++} \to 3l^+ + 3\nu$ may happen.

The models with $\beta = -\sqrt{3}$ cannot be embedded, at least in a straightforward way, in a grand unified theory (GUT) since the existence of exotic quarks implies that new extra exotic quarks has to be added in a supermultiplet. However, who said that the road to GUT theories will be straightforward? Moreover, these model has a Landau-like pole that make them non-perturbative at a few TeVs [17], hence at this scale new degrees of freedom could arise.

After we have finished this work we saw that in Ref. [18] similar left-right symmetric extensions of 3-3-1 models were considered. Those authors consider a general analysis of the models in term of the β parameter in the definition of the electric charge operator. In our case we have considered $\beta = -\sqrt{3}, -1/\sqrt{3}$ each one with two different representation content. It is worthing to note that some characteristics do not depend on the value of β. For instance, in the model with $\beta = -\sqrt{3}$ we have two possibilities in the lepton sector: i) $\psi_l = (\nu_l\, l\, l^c)_L$, $\psi_l = (\nu_l\, E\, E^c)_R$; or, ii) $\psi_l = (\nu_l\, l\, \mathrm{E})_L$, $\psi_l = (\nu_l\, l\, \mathrm{E})_R$. This difference is not trivial, in the first case the flavour lepton numbers is violated because the charged lepton and their charge conjugate are in the same triplet while in the second case we can assign to E the same flavour lepton number of the known charged leptons. This difference implies a different mechanism for generating the lepton masses as can be appreciated by comparing the mass matrices in Eq. (7) with that in Eq. (8). Furthermore, another difference is evident comparing the matrices in Eq. (7) with those in Eq. (17), in particular neutrinos are Majorana, in the former case, and Dirac in the second one. Moreover in the case of Eq. (17) a type-I Dirac seesaw mechanism is implemented. Phenomenology independently of the value of β can be made is some simple situation as, for instance, considering only the Z' of the models [19].

---

[1] R. N. Mohapatra and J. C. Pati, A Natural Left-Right Symmetry, Phys. Rev. D **11**, 2558 (1975).

[2] G. Senjanovic and R. N. Mohapatra, Exact Left-Right Symmetry and Spontaneous Violation

of Parity, Phys. Rev. D **12**, 1502 (1975).

[3] G. Senjanovic, Spontaneous Breakdown of Parity in a Class of Gauge Theories, Nucl. Phys. B **153**, 334 (1979).

[4] R. N. Mohapatra and R. E. Marshak, Local B-L Symmetry of Electroweak Interactions, Majorana Neutrinos and Neutron Oscillations, Phys. Rev. Lett. **44**, 1316 (1980); Erratum: [Phys. Rev. Lett. **44**, 1643 (1980)].

[5] R. N. Mohapatra and G. Senjanovic, Neutrino Mass and Spontaneous Parity Violation, Phys. Rev. Lett. **44**, 912 (1980).

[6] R. N. Mohapatra and G. Senjanovic, Neutrino Masses and Mixings in Gauge Models with Spontaneous Parity Violation, Phys. Rev. D **23**, 165 (1981).

[7] R. A. Diaz, R. Martinez and F. Ochoa, SU(3)(c) x SU(3)(L) x U(1)(X) models for beta arbitrary and families with mirror fermions, Phys. Rev. D **72**, 035018 (2005); [hep-ph/0411263].

[8] A. G. Dias, C. A. de S.Pires and P. S. Rodrigues da Silva, The Left-Right SU(3)(L)xSU(3)(R)xU(1)(X) Model with Light, keV and Heavy Neutrinos, Phys. Rev. D **82**, 035013 (2010) [arXiv:1003.3260 [hep-ph]].

[9] F. Pisano and V. Pleitez, $SU(3)_L \otimes U(1)_N$ *model for electroweak interactions*, Phys. Rev. D **46**, 410 (1992); [hep-ph/9206242].

[10] P. H. Frampton, Chiral dilepton model and the flavor question, Phys. Rev. Lett. **69**, 2889 (1992).

[11] R. Foot, O. F. Hernandez, F. Pisano and V. Pleitez, Lepton masses in an SU(3)-L x U(1)-N gauge model, Phys. Rev. D **47**, 4158 (1993); [hep-ph/9207264].

[12] V. Pleitez and M. D. Tonasse, Heavy charged leptons in an SU(3)-L x U(1)-N model, Phys. Rev. D **48**, 2353 (1993); [hep-ph/9301232].

[13] J. C. Montero, F. Pisano and V. Pleitez, Neutral currents and GIM mechanism in SU(3)-L x U(1)-N models for electroweak interactions, Phys. Rev. D **47**, 2918 (1993); [hep-ph/9212271].

[14] R. Foot, H. N. Long and T. A. Tran, $SU(3)_L \otimes U(1)_N$ and $SU(4)_L \otimes U(1)_N$ gauge models with right-handed neutrinos, Phys. Rev. D **50**, no. 1, R34 (1994); [hep-ph/9402243].

[15] M. Singer, J. W. F. Valle and J. Schechter, Canonical Neutral Current Predictions From the Weak Electromagnetic Gauge Group SU(3) X $u(1)$, Phys. Rev. D **22**, 738 (1980).

[16] J. W. F. Valle and M. Singer, Lepton Number Violation With Quasi Dirac Neutrinos, Phys. Rev. D **28**, 540 (1983).

[17] A. G. Dias, R. Martinez and V. Pleitez, Concerning the Landau pole in 3-3-1 models, Eur. Phys. J. C **39**, 101 (2005); [hep-ph/0407141].

[18] M. Reig, J. W. F. Valle and C. A. Vaquera-Araujo, Unifying left-right symmetry and 331 electroweak theories, arXiv:1611.02066 [hep-ph].

[19] A. J. Buras, F. De Fazio and J. Girrbach-Noe, Z-Z' mixing and Z-mediated FCNCs in $SU(3)_C \times SU(3)_L \times U(1)_X$ models JHEP **1408**, 039 (2014); [arXiv:1405.3850 [hep-ph]].